

The meiobenthos of the Molloy Deep (5600 m), Fram Strait, Arctic Ocean

T. SOLTWEDEL, M. MILJUTINA, V. MOKIEVSKY, D. THISTLE, K. VOPEL - 1-13 - [Abstract](#) - [rokdownload menuitem="441" downloaditem="489" direct_download="true"]PDF[/rokdownload]

Vegetation changes from les Llaunes natural reserve (NE Iberian Peninsula) after seventeen years of management: evaluation by means of GIS

J. GESTI, L. VILAR - 15-25 - [Abstract](#) - [rokdownload menuitem="442" downloaditem="465" direct_download="true"]PDF[/rokdownload]

Foraging behavior of the Corsican nuthatch (*Sitta whiteheadi*) in the Corsican pine (*Pinus nigra laricio*)

)
La quête alimentaire de la Sittelle corse (*Sitta whiteheadi*) dans les Pins laricio (*Pinus nigra laricio*)

)
P. VILLARD, S. BICHELBERGER, J.-F. SEGUIN, J.-C. THIBAUT - 27-32 - [Abstract](#) - [rokdownload menuitem="443" downloaditem="466" direct_download="true"]PDF[/rokdownload]

Birds and climate change / Avifaune et changement climatique

F. ARCHAUX - 33-41 - [Abstract](#) - [rokdownload menuitem="444" downloaditem="467" direct_download="true"]PDF[/rokdownload]

Records of *Ceratoperidinium* Margalef (Dinophyceae) from the Mediterranean sea

F. GOMEZ, M. ABOUD-ABI SAAB - 43-46 - [Abstract](#) - [rokdownload menuitem="445" downloaditem="468" direct_download="true"]PDF [rokdownload]

Free-living nematodes communities, indicators of anthropogenic degree in the Bou Ghrara lagoon (Tunisia)

Les peuplements de Nématodes libres, indicateurs du degré d'anthropisation de la lagune de Bou Ghrara (Tunisie)

E. MAHMOUDI, H. BEYREM, P. AISSA - 47-59 - [Abstract](#) - [rokdownload menuitem="446" downloaditem="469" direct_download="true"]PDF[/rokdownload]

A clarification on "gallertoid" and "benthoblastaea-bilaterogastraea" as preferable models for metazoan evolution

T. SYED - [rokdownload menuitem="447" downloaditem="470" direct_download="true"]PDF[/rokdownload]